

Online Supplementary Material to:

The effect of limb amputation on standing weight distribution in the remaining three limbs in dogs

Grayson Lee Cole; Darryl Millis

Vet Comp Orthop Traumatol 2017; 30:

<https://doi.org/10.3415/VCOT-16-05-0075>

Appendix Table 1: Load distribution in the remaining limbs post limb amputation

Patient	Age (yrs)	Sex	Weight (kg)	Breed	Amputated	Contralateral forelimb	Ipsilateral hindlimb	Contralateral hindlimb	Limb amputated
1	1.5	MC	16	Doberman Pinscher	0	45	24	31	Left forelimb
2	4	MC	30	American Staffordshire Terrier	0	52	18	30	Right forelimb
3	0.5	FS	11	American Staffordshire Terrier	0	57	25	18	Left forelimb
4	1	F	19	Samoyed	0	55	24	21	Right forelimb
5	9	MC	25.7	Shetland Sheepdog	0	47	26	27	Right forelimb
6	3	FS	13.4	Mixed breed	0	29	40	31	Right forelimb
7	9	MC	14.5	Keeshond	0	53	24	23	Right forelimb
8	11	FS	23.2	German Shorthaired Pointer	0	47	27	26	Left forelimb
9	1.5	F	12.7	Mixed breed	0	51	23	26	Left forelimb
10	1	MC	23.6	Mixed breed	0	49	25	26	Left forelimb
11	13	FS	14.5	Mixed breed	36	38	26	0	Right hindlimb
12	7	FS	25.5	Mixed breed	37	33	30	0	Left hindlimb
13	9	MC	37.7	Golden Retriever	35	43	22	0	Right hindlimb
14	5	FS	22	Catahoula Leopard Dog	42	30	28	0	Left hindlimb
15	10	FS	30.1	Flat-Coated Retriever	34	33	33	0	Left hindlimb
16	8	FS	12.7	Mixed breed	24	52	24	0	Left hindlimb
17	7	FS	9.5	Shetland Sheepdog	27	42	31	0	Left hindlimb
18	1	FS	21.5	Mixed breed	29	35	36	0	Left hindlimb
19	5	MC	25.3	American Staffordshire Terrier	30	49	21	0	Left hindlimb
20	9	FS	25	Great Pyrenees Dog	30	41	29	0	Left hindlimb